

"Made available under NASA sponsorship
in the interest of early and wide
semination of Earth Resources Survey
Program information and without liability
for any use made thereof."

E72-10214
CR-129184

TYPE I REPORT

TITLE: STRUCTURAL AND LITHOLOGIC STUDY OF NORTHERN CALIFORNIA
COAST RANGES AND SACRAMENTO VALLEY, CALIFORNIA

PRINCIPAL INVESTIGATOR: Ernest I. Rich
School of Earth Sciences
Stanford University
Stanford, California 94305

PROPOSAL NO.: SR 042

GSFC IDENTIFICATION NO.: UN 217

The first ERTS-1 data was received by the Principal Investigator
on October 23, 1972.

Preliminary analysis of the data received has disclosed two po-
tentially important northwest-trending systems of linear features with-
in the Northern California Coast Ranges and a third system, which trends
northeast, can be traced with great uncertainty across the alluviated
part of the Sacramento Valley and into the foothills of the Sierra
Nevada. These linear features may represent fault systems or zones of
shearing because topographic offset and stratigraphic disruption can be
seen along one or two of the lineations. One of the systems of linear
features is parallel or subparallel with the San Andreas fault and is
confined to the Coastal belt (here referred to as the Coastal System).
Another set, which acutely joins but does not appear to transect the
Coastal System, is confined to the central part of the Coast Ranges.
This system, subsequently referred to as the Central System, is less
well defined but appears to be the chief influence on the structural
and topographic grain of the terrain. The third set of linear features,
the Valley System, trends northeasterly, about normal to the Central
system. Stratigraphic offset and/or disruption of the Late Mesozoic
sedimentary rocks along the western margin of the Sacramento Valley and
in the western foothills of the Sierra Nevada can be seen on the imagery.
Northeasterly trending linear elements, subparallel with the Valley
System, can be detected within the alluviated part of the Sacramento
Valley. These elements may reflect the continuation of the Valley System
within the bedrock that floors the Sacramento Valley. The Valley System
has not been recognized heretofore. Because of the limited imagery
coverage received so far, the extent of these linear features or their
tectonic significance cannot be ascertained.

One, and perhaps two, stratigraphic unconformities within the
Late Mesozoic sedimentary rocks were detected during preliminary exami-
nation of the imagery; however, more careful analysis of subsequent
imagery and field examination is required to verify this preliminary
interpretation.

(E72-10214) STRUCTURAL AND LITHOLOGIC
STUDY OF NORTHERN CALIFORNIA COAST RANGES
AND SACRAMENTO VALLEY, CALIFORNIA E.I.
Rich (Stanford Univ.) [1972] 2 p CSCI 08G
N73-12332
G3/13
Unclas 00214

Analysis of U-2 imagery supplied by NASA Ames Research Center has been completed and photogeologic criteria for the various rock types has been established for the different spectral bands of ERTS. Because of the seasonal difference between the U-2 imagery and the ERTS-1 imagery presently at hand, no direct correlation of photogeologic criteria is possible. However, subtle tonal and topographic differences within areas familiar to the Principal Investigator, suggest that gross rock type differences may be detected on ERTS-1 imagery.